

2004 Annual Report
of
The South Australian Cerebral Palsy Register
(part of The South Australian Birth Defects Register)

**Children born 1993 to 1999 with cerebral palsy
notified to the Register by 31st December 2004**

August 2005

The South Australian Cerebral Palsy Register

Location

Women's and Children's Hospital
Level 8 Rieger Building
72 King William Road
North Adelaide SA 5006
Telephone: (08) 8161 7242
Fax: (08) 8161 6088
Email: cprester@mail.wch.sa.gov.au
www.wch.sa.gov.au/services/az/divisions/labs/geneticmed/cerebral_palsy.html

Management Committee

Prof Eric Haan	Clinical Geneticist. Head, SA Clinical Genetics Service
Dr Annabelle Chan	Public Health Physician
Dr Peter Flett	Paediatric Rehabilitation Specialist. Director, Paediatric Rehabilitation
Dr Deirdre White	Developmental Paediatrician.
Ms Terry Lyons	Clinical Manager Occupational Therapy Novita Children's Service
Ms Phillipa van Essen	Senior Project Officer. SA Birth Defects Register
Ms Kylie Tugaraza	Coordinator, Cerebral Palsy Register

Steering Committee

Women's and Children's Hospital

Dr Peter Flett Paediatric Rehabilitation Specialist

Neonatal Follow-Up Program: Women's and Children's Hospital

A/Prof Ross Haslam Neonatal Paediatrician

Neonatal Follow-up Program: Flinders Medical Centre

A/Prof Peter Marshall Neonatal Paediatrician

South Australian Birth Defects Register (SABDR)

Prof Eric Haan Clinical Geneticist

Collaborating Organisations

Novita Children's Services
Flinders Medical Centre
Women's and Children's Hospital

Funding

The Register is funded by the Women's and Children's Hospital, with additional support provided by Novita Children's Services of South Australia and the Community Accommodation and Respite Agency (CARA).

The South Australian Cerebral Palsy Register – Annual Report 2004

Contents

Foreword	4
Management Committee, South Australia Cerebral Palsy Register	5
Summary	6
Thanks to notifiers / Acknowledgments	7
What is cerebral palsy?	8
Funding / Administration / Ethical approval / Legislative framework /Aims of the Register	9
How the Register works	10
Register activities	11
Prevalence of cerebral palsy 1993-99	12
Notifiers to the Register	13
Type of cerebral palsy	14
Gestational age at birth	15
Birth weight	16
Level of motor function	17
Co-morbidities	18
Presentations / Conferences	21
Publications	22
References	27

Tables, Figures and Appendices

Tables & Figures

All notified children with cerebral palsy born in South Australia

Table 1	SA children with cerebral palsy ascertained by the Register 1993-99	12
Table 2	Cerebral palsy ascertainment and prevalence 1993-99	12
Table 3	Notifiers to the Register 1993-99	13
Table 4	Type of cerebral palsy 1993-99	14
Table 5	Cerebral palsy by gestational age at birth 1993-99	15
Table 6	Cerebral palsy by birth weight range 1993-99	16
Figure 1	Notifiers to the Register 1993-99	13
Figure 2	Type of cerebral palsy 1993-99	14
Figure 3	Cerebral palsy by gestational age at birth 1993-99	15
Figure 4	Cerebral palsy by birth weight 1993-99	16

Children with cerebral palsy born in South Australia and assessed at five years of age

Table 7	Level of motor function 1993-99	17
Table 8	Intellectual ability 1993-99	18
Table 9	Visual problems 1993-99	19
Table 10	Hearing problems 1993-99	19
Table 11	Types of cerebral palsy and associated disabilities	20
Figure 5	Level of motor function 1993-99	17
Figure 6	Intellectual ability 1993-99	18

Appendices

Appendix 1	Gross motor function classification system in cerebral palsy	23
Appendix 2	South Australian Cerebral Palsy Register Family Information Sheet	25

Foreword

When the South Australian Cerebral Palsy Register was first set up in 1997 it had several important aims, which included obtaining information to help in the planning of the services and needs of those with cerebral palsy and their families. It was also envisaged at that time that the enlarging database might increasingly be useful for epidemiological research into the causation of cerebral palsy. Indeed data from the SA register and the registers in Western Australia and Victoria have since confirmed strong associations between cerebral palsy and prematurity, intrauterine growth restriction, intrauterine infection and multiple pregnancies.


However, the cerebral palsy register is now proving to be a wonderful research tool in a way that perhaps was not envisaged in 1997. Using the extensive data in the register, the clinical details of hundreds of cases can be linked with new molecular biological tests, which are helping to understand the fetal origins of cerebral palsy.

Until recently it had been a popular but mistaken belief that cerebral palsy frequently began in labour or at birth because of low oxygen levels or trauma from the delivery. There is now increasing evidence that the “fetal distress” or poor condition after birth, of babies who develop cerebral palsy, may often be a reflection of a longer standing pathological process or genetic vulnerability. This may explain why no modern obstetric interventions in labour such as electronic fetal heart rate monitoring or a large increase in caesarean section deliveries have changed the cerebral palsy rates. Indeed currently there is no obstetric clinical policy before or during labour that has been shown to reduce the risk of cerebral palsy.

With the help of the SA Cerebral Palsy Register it has been possible to select (without access to the identity of the child) stored blood spots from these children and from control babies who did not develop cerebral palsy. Using a tiny (1 mm) punch from a dried blood spot it has been possible through DNA analyses to identify genetic aberrations that increase the risk in the fetus to infection (cytokine polymorphisms) and clotting (hereditary thrombophilias). On the same blood spot, evidence of viral exposure around pregnancy can be identified. This novel ongoing research has identified important associations between cerebral palsy and an increase in the above infective and hereditary risk factors. Current research is investigating the interaction of genetic and environmental factors such as growth restriction that may potentiate the risk of future cerebral palsy. This is leading to a better understanding of the multifactorial causes of cerebral palsy and in time will lead to the testing of interventions before, during and soon after pregnancy, to reduce the risk of cerebral palsy.

Although cerebral palsy rates have not reduced in the last 40 years the recent genetic breakthroughs in research into cerebral palsy causation are likely to lead, in the next decade, to preventative treatments and policies that should decrease cerebral palsy rates. It will be a pleasure to see the SA Cerebral Palsy Register being used to monitor and validate these possible reductions, which were facilitated by the very existence of the register itself.

I offer my congratulations to the management and steering committees and to the Head of the SA Cerebral Palsy Register, Professor Eric Haan who, this year, deservedly was awarded the Order of Australia.



**Professor Alastair MacLennan, The University of Adelaide,
Chairman of the South Australian Cerebral Palsy Research Group.**



Management Committee, South Australian Cerebral Palsy Register



Management Committee South Australian Cerebral Palsy Register

Front Row: Ms Heather Scott, Dr Annabelle Chan, Dr Deirdre White,

Back Row: Dr Peter Flett, Ms Kylie Tungaraza, Prof Eric Haan,

Absent: Terry Lyons, Phillipa van Essen.

Summary

The South Australian Cerebral Palsy Register is a population-based collection of information on children with cerebral palsy born in South Australia in a population with an average of 19075 live births per year (1993-1999).

This report presents information for children with cerebral palsy born in the years 1993-1999.

The term cerebral palsy describes a group of permanent and non-progressive disorders of movement and posture that manifest early in life and result from a defect or lesion of the immature brain. Individuals with cerebral palsy have life-long motor disabilities, frequently associated with intellectual disability, epilepsy and visual and hearing impairment. It is the most common chronic motor disability of childhood and places a large emotional and financial burden on those affected and their families.

The types of cerebral palsy are based on clinical features and comprise spastic quadriplegia, spastic triplegia, spastic diplegia, spastic hemiplegia, monoplegia, ataxia and dyskinesia.

Children with cerebral palsy are ascertained through notifications to the South Australian Birth Defects Register from Novita Children's Services, rehabilitation specialists, other paediatricians and the Neonatal Long Term Follow-up Programs of Flinders Medical Centre and Women's and Children's Hospital.

Around five years of age, a comprehensive medical history is obtained and the children have a formal clinical assessment to ensure that the diagnosis of cerebral palsy is correct, to document the type and severity of the disorder, and to define the disabilities experienced by the child.

Parental consent is obtained for children to participate in the clinical assessment at 5 years of age.

For the years 1993-1999, 285 children with cerebral palsy were ascertained. At 31st December 2004, 68% of these children had had their comprehensive clinical assessment at around five years of age.

The maximum and minimum prevalence rate of cerebral palsy between 1993-1999 per 1000 live births is presented. Minimum rate represents the cases confirmed by examination at five years, and the maximum rate reflects the total ascertained cases. Between 1993-1999 the minimum rate of cerebral palsy was 1.46 per 1000 live births, and the maximum rate of cerebral palsy 2.13 per 1000 live births.

Paediatric rehabilitation specialists notified 39% of cases, Novita Children's Services 31%, the South Australian Birth Defects Register 21%, other specialists 4%, and Other sources 4%.

Hemiplegia (33%) and diplegia (32%) were the most common forms of cerebral palsy, followed by quadriplegia (21%), triplegia (5%) and ataxia (3%). Dyskinetic athetoid (2%), monoplegia (1%) and dyskinetic dystonic (<1%) were less common forms of the disorder. In 3% of cases, the type of cerebral palsy was unknown.

48.1% of affected children were born at term (≥ 37 weeks), 51.9% were born prematurely, and none had an unknown gestation.

The association between the risk of cerebral palsy and prematurity is exemplified by a prevalence of 87.7 cases per 1000 live births at 23-27 weeks gestation compared to 1.0 case per 1000 live births at term (≥ 37 weeks).

50.9% of affected children had a birth weight ≥ 2500 grams, 17.9% weighed 1500-2499 grams, 14% weighed 1000-1499 grams, and 17.2% weighed <1000 grams.

The prevalence of cerebral palsy was 88.4 cases per 1000 among live births with birth weights in the 500-999 g range, but only 0.8 per 1000 among live births with a birth weight in the 3500-3999 g range.

66% of affected children had relatively mild cerebral palsy (Levels I and II) while 33% had more severely affected gross motor function (Levels III-V); and 1% of cases had an unknown level of motor function.

39.5% of children with cerebral palsy born between 1993 and 1999 had impaired intellectual ability. 37% had impaired vision and 14% had impaired hearing.

Thanks to Notifiers / Acknowledgments

We wish to thank all notifiers to the register, including paediatricians, staff of Novita Children's Services of South Australia and the families of children with cerebral palsy. In particular we would like to thank the staff of the Department of Child and Adolescent Development, Neurology and Rehabilitation of the Women's and Children's Hospital. We acknowledge the ongoing assistance of the staff of the Department of Medical Records at Flinders Medical Centre, The Queen Elizabeth Hospital, Lyell McEwin Health Services and Women's and Children's Hospital. Special thanks must also go to staff of the Pregnancy Outcome Statistics Unit of the Epidemiology Branch, Department of Health for their invaluable assistance.

Acknowledgments

The Register gratefully acknowledges funding from the Community Accommodation and Respite Agency (CARA), which allowed the Register to upgrade its computing in 2004.

We would like to thank all those people and organisations who have contributed to the South Australian Cerebral Palsy Register during its planning stage and since its inception; their ongoing support and practical contribution is much appreciated.

Special thanks are due to:

- Novita Children's Services, formerly Crippled Children's Association of South Australia (CCA)
- The Community Accommodation and Respite Agency (CARA)
- Paediatricians from specialist centres and in private practice
- Staff of the Department of Child and Adolescent Development, Neurology and Rehabilitation of the Women's and Children's Hospital
- Long Term Follow-Up Program coordinators from the Women's and Children's Hospital and Flinders Medical Centre
- Staff of the South Australian Clinical Genetics Service, Women's and Children's Hospital.

While there are many who have contributed to the Cerebral Palsy Register, there would be no Register without the kind cooperation of the families of affected children. These families live with the effects of cerebral palsy on a daily basis. Once again the Register owes them a debt of gratitude for allowing us to enter their lives and for their insightful comments over the year.

What is Cerebral Palsy?

Cerebral palsy is a term of convenience applied to a group of motor disorders of central origin defined by a clinical description (Badawi et al, 1999) ¹. It covers a range of cerebral disorders that result in childhood motor impairment. The impairment must stem from non-progressive malfunction of the brain (rather than the spinal cord or muscles).

Cerebral palsy affects an individual's ability to control movement and posture. Palsy is an expression used to describe paralysis. A more accurate description of the muscle symptoms might be weakness (paresis) and an inability to make voluntary movements and to suppress involuntary ones.

Unlike many other disorders that affect the motor system, cerebral palsy is not progressive - it does not get worse with time.

In fact, family nurturing, therapy and education can result in improvements in functional outcomes and quality of life.

Approximately 2 in every 1000 children born in South Australia have cerebral palsy, which equates to approximately 38 newly diagnosed children each year in 1993-1999. The incidence of cerebral palsy has not decreased, despite improvements in ante-, intra- and post-natal care.

Children affected by cerebral palsy will have life-long disabilities and continue to represent a significant proportion of the children and young adults with disabilities in the community. Severity can vary greatly, from minor awkwardness to severe multiple disabilities

Cerebral palsy can be associated with other disorders, such as epilepsy, learning difficulties, and problems with sight and hearing. There can be difficulties with communication, resulting from multiple disabilities, including problems controlling the speech muscles, intellectual disabilities and poor hearing.

Orthopaedic complications are common, often requiring ongoing care. Most children with cerebral palsy survive to adulthood, although severely affected children may have a reduced life expectancy.

Whilst there is no single cause for cerebral palsy, we do know that there are several perinatal factors which are associated, both individually and in combination, with an increased risk of cerebral palsy. These include:

- Prematurity and very low birth weight
- Maternal infections, such as cytomegalovirus (CMV), group B streptococcus and rubella
- Feto-maternal haemorrhage
- Hypoxia during labour and delivery
- The use of certain drugs during pregnancy, such as cocaine
- Excessive alcohol intake during pregnancy
- Metabolic problems in the newborn period, such as severe jaundice and hypoglycaemia.

Recent research using blood specimens from babies obtained soon after birth, has demonstrated an association between markers of prenatal inflammation and certain coagulation abnormalities, and the subsequent development of cerebral palsy. (Nelson et al, 1998) ².

There are also events occurring in early childhood (before age 2 years) which may result in "acquired" cerebral palsy. These include:

- Near drowning
- Brain hypoxia from any cause
- Some brain tumours
- Head injuries: accidental and non-accidental
- Severe infection involving the brain, such as meningitis or encephalitis
- Cerebrovascular accidents
- Non accidental injuries
- Complications of inborn errors of metabolism.

Funding / Administration / Ethical approval / Legislative framework

Funding

The Register is funded by the Women's and Children's Hospital, with additional funding from Novita Children's Services (formerly Crippled Children's Association CCA), Community Accommodation & Respite Agency (CARA) and the Rotary Club of Morialta supported the establishment phase of the Register. The Register gratefully acknowledges funding from the Community Accommodation and Respite Agency (CARA), in 2004 that allowed the Register to upgrade its computing.

Administration

The Register is based at the Women's and Children's Hospital, and is located administratively within the South Australian Clinical Genetics Service. Its location enables it to draw on the experience and expertise of the Department of Child and Adolescent Development, Neurology and Rehabilitation of the Women's and Children's Hospital. The Register's Management Committee is drawn from members of these departments and Novita Children's Services.

Ethical approval

The establishment of the Cerebral Palsy Register was approved by the Human Research Ethics Committees of the Women's and

Children's Hospital, the North Western Adelaide Health Service and Flinders Medical Centre. Continuing ethical oversight of the Register is carried out by the Human Research Ethics Committee of the Women's and Children's Hospital.

Legislative framework

The Cerebral Palsy Register is part of the SA Birth Defects Register and operates under the provisions of the South Australian Health Commission (Pregnancy Outcome Statistics) Regulations of 1999. This Regulation requires the notification of all congenital abnormalities diagnosed before the child's fifth birthday. Section 64d of the South Australian Health Commission Act, 1976, Amendment Act, 1989 allows the Register to carry out research into cerebral palsy while making provision for maintaining the confidentiality of participants.

Aims of the Register

- To determine and monitor the prevalence of cerebral palsy in South Australia
- To gather information about affected children that may provide clues to the causes of cerebral palsy
- To document the severity and range of disabilities experienced by children with cerebral palsy
- To use the information collected to plan facilities for affected children
- To act as a source of information about cerebral palsy, for both families and the community
- To improve community and professional awareness of cerebral palsy, including its causes and outcomes
- To provide a resource for research into cerebral palsy
- To contribute to mortality and morbidity studies of cerebral palsy.

How the Register Works

Ascertainment

Notifications of children with cerebral palsy come from a variety of sources, including Novita Children's Services, paediatricians, physiotherapists, neurologists and occupational therapists. The main sources of notification come from the Women's and Children's Hospital/ Novita Paediatric Rehabilitation Specialists. The Register encourages multiple notification as a means of ensuring complete ascertainment of affected children.

All notifications are received by the SA Birth Defects Register. Once notified, the Register approaches families through a clinician known to them, who will be one of the physicians working with the Register or the child's paediatrician. Information about the Register and its aims is sent to the managing clinician and an invitation to become involved in the Register is extended to the family. The managing clinician is also sent a package containing a family information leaflet and a 'consent to contact' form. Once the clinician has communicated with the family, the package can be forwarded to them. The family can then respond directly to the Register.

The 'consent to contact' form gives the Register permission to contact the family, allowing the Register coordinator to give the family detailed information about the Register's aims and functioning. Families are free to refuse participation in the Register. Those who agree to participate may, at any time, ask to be withdrawn from the Register with the knowledge that this will not affect any aspect of the child's medical care. Children included on the Register are given a full clinical assessment at five years of age, performed by a paediatrician working with the Register. By then, it is clear whether or not cerebral palsy is present, and the type and severity can be determined more accurately. Disabilities in addition to the motor disability can also be documented.

The purpose of the assessment is to collect data on the type of cerebral palsy, its severity and any associated disabilities. Information about learning problems, epilepsy, hearing, vision and the use of mobility aids is also collected. A brief family history is obtained to determine if cerebral palsy, intellectual disability or epilepsy has occurred within the immediate or extended family in the past. The method of the child's conception is documented. Information relating to the perinatal period is also collected.

Consents are obtained from families to allow the Register Coordinator to access information from a variety of sources. The Register works closely with the Pregnancy Outcome Statistics Unit, medical records departments and clinicians in order to ensure accurate and complete data for each child.

Data Storage

The collected data are stored on computer and are protected by using unique identifier codes to maintain participant confidentiality. All information and participant files are backed up regularly to prevent data loss.

Confidentiality

The Register does not release identified information about a participant without the written consent of his or her parents or guardian.

Inclusion and Exclusion Criteria

The Register includes children who have a motor impairment, manifested early in life, which is the result of static cerebral pathology. The Register collects post-neonatally acquired cases of cerebral palsy (after the first month and before age 2 years).

The following disorders are excluded: neurodegenerative conditions, neuromuscular disorders, neural tube defects, tumours, hypotonia occurring in isolation or with intellectual disability, many genetic syndromes and most inborn errors of metabolism. There are many conditions where a decision about inclusion or exclusion can be difficult, and guidance is provided by Badawi et al (1998).

Register activities 2003/2005

During 2003 / 2005, Register staff have:

- Continued to refine the data set, database and systems for analysis and interpretation of data
- Continued to participate in the South Australian Cerebral Palsy Research Group, which is investigating whether variations in genes for clotting factors and perinatal infection are associated with cerebral palsy.
- Received general queries about setting up a cerebral palsy register from representatives in other Australian States and Territories requests for information on consent forms and information pamphlets and legislation.
- Received requests for information from the community and tertiary students who required statistics on the prevalence of cerebral palsy; types of cerebral palsy and associated disabilities in South Australia.
- Received requests for information on general information of cerebral palsy for cerebral palsy week from the Department of Disabilities Family and Community.
- Continued to collaborate with staff from the Department of Child and Adolescent Development, Neurology and Rehabilitation of the Women's and Children's Hospital as well as staff from the Repatriation General Hospital on a project investigating whether Botox injections can improve upper limb function in children with hemiplegic cerebral palsy.
- Attended the meeting of the Australian National Cerebral Palsy Register's State and Territory Representatives on Thursday 3 April 2003 in Sydney to discuss standards and methodology for the National Cerebral Palsy Database, including signing a memorandum of understanding on how the data will be utilised.
- Attended the meeting of the Australian National Cerebral Palsy Register's State and Territory Representatives on Monday 16 and Tuesday 17 May 2005 in Perth to discuss national classification issues and finalise the minimum data set.

Prevalence of cerebral palsy 1993 - 1999

Tables 1-6 and Figures 1-4 present data about children with cerebral palsy notified to the South Australian Birth Defects Register.

Table 1

SA children with cerebral palsy ascertained by the Register 1993-99

Group	1993	1994	1995	1996	1997	1998	1999	1993-99	
	No.	No.	No.	No.	No	No	No	No.	(%)
Assessment completed at 5 years									
- Including clinical examination	31	31	42	20	40	20	11	195	(68)
- Medical records and reports only [#]	4	8	11	8	14	20	16	81	(28)
Deceased prior to 5 years	1	4	0	1	0	3	0	9	(3)
Total*	36	43	53	29	54	43	27	285	(100)

[#] Two families had moved interstate.

* Does not include interstate or overseas births:- two in 1993, four in 1994, six in 1995, seven in 1996, five in 1997, four in 1998 and four in 1999.

Two hundred and eighty five children, born in South Australia (SA) in 1993 – 1999, had been notified to the Register by 31st December 2004. Over this same period, the Register also received 32 notifications of cases born outside SA. A comprehensive clinical assessment at the age of five years has been completed for 195 (68%) of the 285 children (see Table 1).

Experience has shown that some of the notified cases will not have cerebral palsy when assessed at five years of age; for this reason minimum and maximum rates has been calculated for the years 1993 – 99 (Table 2). Minimum rate represents confirmed cases and the maximum rate reflects total ascertained cases.

Table 2

Cerebral palsy ascertainment and prevalence 1993-99

Year of birth	Live births	Ascertained cases	Confirmed cases* (%)	Rate per 1000 (min, max) [#]
1993	19846	36	31 (86%)	1.56, 1.81
1994	19673	43	31 (72%)	1.58, 2.19
1995	19472	53	42 (79%)	2.16, 2.72
1996	18979	29	20 (69%)	1.05, 1.53
1997	18535	54	40 (74%)	2.16, 2.91
1998	18613	43	20 (47%)	1.07, 2.31
1999	18404	27	11 (41%)	0.60, 1.47
1993-99	133522	285	195 (68%)	1.46, 2.13

*Have completed comprehensive clinical assessment at 5 years of age.

[#] Rate per 1000 live births. Minimum rate represents confirmed cases and maximum rate reflects total ascertained cases.

Notifiers to the Register


The sources of notification of children with cerebral palsy born in SA between 1993 and 1999 are shown in Table 3. Over this period, 92% of the notifications came from three main sources, Rehabilitation Specialists, Novita Children's Services and the South Australian Birth Defects Register.

Table 3
Notifiers to the Register 1993-99

Notifier*	1993	1994	1995	1996	1997	1998	1999	1993-99	
	No.	No.	No.	No.	No.	No.	No.	No.	(%)
Rehabilitation specialists	12	5	16	15	30	24	10	112	(39.3)
Other specialists	2	0	0	2	4	0	2	10	(3.5)
Novita Children's Services	14	33	26	9	2	5	0	89	(31.2)
SA Birth Defects Register	5	3	7	3	17	13	12	60	(21.1)
Other sources	3	2	4	0	1	1	0	11	(3.9)
Unknown	0	0	0	0	0	0	3	3	(1.1)
Total	36	43	53	29	54	43	27	285	(100)

*The Register encourages multiple notifications.

Figure 1
Notifiers to the Register 1993-99



Type of cerebral palsy


The different types of cerebral palsy for children born in SA between 1993 and 1999 are shown in Table 4. The most common forms of cerebral palsy were hemiplegia, diplegia and quadriplegia accounting for 86% of all cases.

Table 4
Type of cerebral palsy 1993-99

Type of cerebral palsy	1993 No.	1994 No.	1995 No.	1996 No.	1997 No.	1998 No.	1999 No.	1993-99 No.	1993-99 (%)
Quadriplegia *	7	9	12	3	8	11	9	59	(20.7)
Triplesia	1	2	3	4	3	1	0	14	(4.9)
Hemiplegia	10	16	18	9	19	11	11	94	(33.0)
Diplegia#	16	16	14	11	17	12	6	92	(32.3)
Monoplegia	0	0	0	0	0	2	0	2	(0.7)
Ataxia	1	0	1	0	1	4	1	8	(2.8)
Dyskinetic athetoid	1	0	3	1	1	0	0	6	(2.1)
Dyskinetic dystonic	0	0	0	0	1	0	0	1	(0.4)
Hypotonia	0	0	1	0	0	0	0	1	(0.4)
Unknown	0	0	1	1	4	2	0	8	(2.8)
Total	36	43	53	29	54	43	27	285	(100)

Figure 2

Type of cerebral palsy 1993-99



* Spasticity in four limbs: Upper limbs greater than or equal to lower limbs

Spasticity in four limbs: Lower limbs greater than upper limbs

Gestational age at birth

Gestational ages at birth for cases of cerebral palsy born in SA between 1993-99 are shown in Table 5. 48% of cases were born at term (≥ 37 weeks).

The rate of cerebral palsy by gestational age at birth 1993-99 demonstrated in Table 5 and Figure 3 illustrates the association between cerebral palsy and prematurity (<37 weeks), especially gestations ≤ 31 weeks.

Table 5
Cerebral palsy by gestational age at birth 1993-99

Gestation* Weeks	1993	1994	1995	1996	1997	1998	1999	1993-99		1993 – 99
	No.	No.	No.	No.	No.	No.	No.	No.	(%)	Rate [#]
23 [^] - 27	2	6	6	8	12	14	2	50	(17.5)	87.7
28 - 31	9	7	13	3	5	6	7	50	(17.5)	41.3
32 - 36	6	8	9	3	9	7	6	48	(16.9)	5.2
37 - 41	19	22	24	14	28	16	12	135	(47.4)	1.0
42+	0	0	1	1	0	0	0	2	(0.7)	0.9
Total	36	43	53	29	54	43	27	285	(100)	2.13


*Gestation at birth (best clinical estimate) in weeks

[^] There were no cases below 23 weeks

[#] Rate per 1000 live births

Figure 3

Cerebral palsy by gestational age at birth 1993-99



Birth weight

Weight ranges at birth from 1993-99 are shown in Table 6 and Figure 4. From the Table it can be seen that there is an extensive spread over the birth weight ranges. The ranges 500-999 g (17.2%) and 3000-3499g (17.5%) were the most common weight ranges for children with cerebral palsy, followed by 2500-2999g (15.1%).


The rates of cerebral palsy per 1000 live births, by birth weight, are provided in Table 6 and Figure 4 for the period 1993-1999. They demonstrate a strong relationship between cerebral palsy and low birth weight. As birth weight increases, the rate of cerebral palsy decreases, from 88.4 per 1000 live births at 500 – 999 g to 0.8 per 1000 livebirths at 3500 – 3999 g.

Table 6
Cerebral palsy by birth weight 1993-99

Birth weight range (grams)	1993	1994	1995	1996	1997	1998	1999	1993-99		1993-1999 Rate [#]
	No.	No.	No.	No.	No.	No.	No.	No. (%)		
500 - 999	8	4	10	4	10	12	1	49	(17.2)	88.4
1000 - 1499	4	8	5	4	7	6	6	40	(14.0)	49.6
1500 - 1999	3	5	7	2	0	6	6	29	(10.2)	16.8
2000 - 2499	3	3	3	3	4	3	3	22	(7.7)	4.1
2500 - 2999	3	5	7	9	8	6	5	43	(15.1)	2.1
3000 - 3499	6	8	10	4	14	6	2	50	(17.5)	1.0
3500 - 3999	7	7	6	3	5	2	2	32	(11.2)	0.8
4000 - 4499	1	2	3	0	6	2	2	16	(5.6)	1.2
4500 - 4999	1	1	2	0	0	0	0	4	(1.4)	1.9
Total	36	43	53	29	54	43	27	285	(100)	2.13

[#] Rate per 1000 live births

Figure 4
Rate of cerebral palsy by birth weight 1993-99



Level of motor function

Tables 7-11 and Figures 5-6 present additional data collected at the 5 year assessment. As only 195 (68%) of notified cases between 1993-1999 had a clinical examination as part of the 5 year assessment, the numbers for each year are less than in the previous section of the report.

Table 7 and Figure 5 show the level of motor function for children with cerebral palsy born 1993 – 1999. The majority of children had less restricted motor function Levels I and II i.e. walking without restriction (45.1%) and walking without assistive devices (21.0%) respectively, while 33% were more severely affected in their motor function Levels III–V, with limitations or severe limitations to self-mobility. (A definition of motor function appears in Appendix 1).


Table 7

Level of motor function 1993-99

Level of function*	1993 No.	1994 No.	1995 No.	1996 No.	1997 No.	1998 No.	1999 No.	1993-99 No. (%)
Level I	9	14	22	10	19	7	7	88 (45.1)
Level II	9	9	3	4	11	3	2	41 (21.0)
Level III	2	5	4	3	1	3	1	19 (9.7)
Level IV	5	1	4	2	5	1	1	19 (9.7)
Level V	6	2	9	1	3	5	0	26 (13.3)
Unknown	0	0	0	0	1	1	0	2 (1.0)
Total	31	31	42	20	40	20	11	195 (100)

Figure 5

Level of motor function * 1993-99



* Definition of motor function appears in Appendix 1.

Co-morbidities


Table 8 and Figure 6 illustrate the level of intellectual ability for children with cerebral palsy born 1993 – 1999. They show that 44.1% of children with cerebral palsy had average or greater intellectual abilities (i.e. an intelligent quotient score of ≥ 90), 11.8% of children had low average (80-89 I.Q.), 21.0% had mild intellectual disability, and 18.5% had moderate to profound intellectual disability. There were 6.7% of children with above average intellectual ability.

Table 8
Intellectual ability 1993-99

Intellectual ability #	1993	1994	1995	1996	1997	1998	1999	1993-99	
	No.	No.	No.	No.	No.	No.	No.	No.	(%)
Above average	0	3	2	0	3	3	2	13	(6.7)
Average	10	8	13	11	19	7	5	73	(37.4)
Low average	5	4	7	2	3	1	1	23	(11.8)
Mild disability	3	10	11	5	7	4	1	41	(21.0)
Moderate disability	7	3	3	0	6	2	0	21	(10.8)
Severe disability	5	2	2	1	2	1	0	13	(6.7)
Profound disability	1	1	0	0	0	0	0	2	(1.0)
Uncertain*	0	0	4	1	0	2	2	9	(4.6)
Total	31	31	42	20	40	20	11	195	(100)

Figure 6

Intellectual ability # 1993-99



#Various scales, appropriate to the child's age and physical abilities, were used to assess intellectual ability.

*Not possible to determine intellectual ability.

Co-morbidities

The frequency of visual problems in children with cerebral palsy born 1993 – 1999 is shown in Table 9. 53.3% of children had normal vision and 37.4% had impaired vision. In 7.7% of cases it was uncertain whether a visual problem was present because of difficulty in assessment eg. due to the level of intellectual disability

Table 9
Visual problems 1993-99

Visual problems*	1993 No.	1994 No.	1995 No.	1996 No.	1997 No.	1998 No.	1999 No.	1993-99 No.	(%)
Normal	14	14	20	17	27	8	4	104	(53.3)
Impaired	13	14	16	3	11	9	7	73	(37.4)
Uncertain	4	1	5	0	2	3	0	15	(7.7)
Unknown	0	2	1	0	0	0	0	3	(1.5)
Total	31	31	42	20	40	20	11	195	(100)

* Visual problems include; strabismus +/- amblyopia, refractive error, cortical blindness, nystagmus and optic nerve abnormalities.

The frequency of hearing problems in children with cerebral palsy born 1993 – 1999 is shown in Table 10. Over 80% of children had normal hearing and less than 15% of cases had impaired hearing. In 2.6% of cases the level of hearing was uncertain and in 1.5% of cases it was unknown.

Table 10
Hearing problems 1993-99

Hearing problems*	1993 No.	1994 No.	1995 No.	1996 No.	1997 No.	1998 No.	1999 No.	1993-99 No.	(%)
Normal	24	23	32	18	33	19	10	159	(81.5)
Impaired	6	6	8	2	4	1	1	28	(14.4)
Uncertain	1	2	1	0	1	0	0	5	(2.6)
Unknown	0	0	1	0	2	0	0	3	(1.5)
Total	31	31	42	20	40	20	11	195	(100)

*Hearing problems include; sensorineural and/or conductive hearing loss determined by audiology.

Co-morbidities

Table 11 reveals the types of cerebral palsy with associated co- morbidities of hearing, vision, epilepsy and intellectual disability. Some differences can be seen between for example the left and right hemiplegic types of cerebral palsy, especially with respect to hearing and vision however these were not found to be statistically significant, $p= 0.86$ and $p= 0.93$ respectively. At least 70% of children with quadriplegia had vision impairment, epilepsy and intellectual disabilities.

Table 11
Types of cerebral palsy and associated disabilities

Type of cerebral palsy	Total in category	Hearing		Vision		Epilepsy		Intellectual disability*	
		No.	(%)	No.	(%)	No.	(%)	No.	(%)
Quadriplegia	30	4	(13)	22	(73)	21	(70)	25	(83)
Triplegia	13	2	(15)	4	(31)	2	(15)	5	(38)
Right hemiplegia	38	7	(18)	12	(32)	10	(26)	18	(47)
Left hemiplegia	29	4	(14)	8	(28)	8	(28)	14	(48)
Diplegia	70	9	(13)	22	(31)	13	(19)	37	(53)
Monoplegia	1	0	(0)	0	(0)	0	(0)	1	(100)
Ataxia	6	1	(17)	3	(50)	2	(33)	3	(50)
Dyskinetic athetoid	6	0	(0)	2	(33)	1	(17)	4	(67)
Dyskinetic dystonic	1	0	(0)	0	(0)	1	(100)	0	(0)
Hypotonia	1	1	(100)	1	(100)	0	(0)	1	(100)
Total (% of total)	195	28	(14)	74	(38)	58	(30)	108	(55)

* Known to have intellectual disability of any degree.

Presentations / Conferences / Publications

Presentations

Flett P. Medical Round presentation on "Stroke from a horse, meningitis, and Peliazeus-Merzbacher disease: what do they have in common?" Women's and Children's Hospital, Adelaide 21st March 2005.

Russo R., Murchland S., Sandelance M., Flett P., Brook P. "Functional outcomes in children injected with botulinum toxin. Improving the upper limb to enhance participation level function in children with spasticity and dystonia", Poster presentation, AusACPDM conference, 18th-20th March 2004.

Flett P. Grand Round lectures on "Botulinum toxin A in cerebral palsy: equalising the race between muscles and bones" to Royal Children's Hospital, Brisbane and to Hervey Bay/Bundaberg Hospitals, Queensland, July/Aug 2002.

Flett P. Invited speaker, 2nd National Symposium on Developmental Paediatrics and Child Neurology, Vellore, South India, October 2002; Four lectures entitled: "Assessment of tone and evaluation of spasticity"; "Cerebral Palsy: overview, management challenges and associated problems"; "Anti-spasticity interventions in cerebral palsy"; "Botulinum toxin A in the management of cerebral palsy".

Flett P. "Beyond the Gait". Workshop convenor and chairperson, Women's and Children's Hospital, Adelaide, August 2001.

Boyd R, Noble I, *Flett P*, Corry I, Graham HK. "An economic evaluation of the use of botulinum toxin A, in the conservative management of equinus in children with cerebral palsy". Presented at the American Academy of Cerebral Palsy and Developmental Medicine (AAPDM) meeting 23rd September 2000, Toronto, Canada.

Conferences

Australian National Cerebral Palsy Register's State and Territory Representatives Conference, Telethon Institute For Child Health Research Perth, 16-17 May 2005 (P. Flett, K. Tungaraza, P. van Essen, H. Scott).

Australian Cerebral Palsy Association national conference, Adelaide, 14th- 15th March 2005. (P. Flett).

American Academy of Cerebral palsy and Developmental Medicine meeting, Los Angeles, Sept-Oct 2004. (P. Flett).

Aus ACPDM scientific meeting, Melbourne, March 2004. (P. Flett).

National Cerebral Palsy Register Meeting, Sydney, 3-4 April 2003 (P. Flett, K. Tungaraza, P. Sharpe, H. Scott).

Paediatric Epilepsy Workshop, Hyatt Regency Adelaide, 3 May 2003 (P. Flett).

AFRM/RACP annual scientific meeting, Hobart, May 2003 (P. Flett).

Gait Analysis Course, Royal Children's Hospital, Melbourne, 25-27 June 2003 (P. Flett).

"Has Clinical Gait Analysis Delivered All That it Promised?" Workshop by Prof Kit Vaughan, Adelaide, 24 July 2003 (P. Flett)

State Branch AFRM Scientific Conference, Hahndorf SA, 25-26 October 2003 (P. Flett).

AusACPDM meeting, "Embracing The Evidence", Sydney, 20th-21st September, 2002 (P. Flett)

2nd National Symposium on Developmental Paediatrics and Child Neurology, Vellore, South India, October 2002 (P. Flett).

Stepping Forward in Cerebral Palsy Management, 14th-15th February 2000, Singapore (P. Flett)..

Australian Faculty of Rehabilitation Medicine, 18th-21st August 2000, Melbourne (P. Flett).

American Academy of Cerebral Palsy and Developmental Medicine (AAPDM), September 20th-23rd 2000, Toronto, Canada (P. Flett).

Presentations / Conferences / Publications

Conferences (*cont.*)

Therapy: "Investing in Independence" Workshop, 24th-26th August 1999, Perth, Western Australia (P. Flett).

Publications utilising data from the Register

Gibson CS, MacLennan AH, Hague WM, Haan E, Priest K, Chan A, Dekker GA. Associations between inherited thrombophilias, gestational age and cerebral palsy. *Am J Obstet Gynecol* (In press).

Gibson CJ, MacLennan AH, Rudski Z, Hague WH, Haan EA, Sharpe P, Priest K, Chan A, and Dekker GA. The prevalence of inherited thrombophilias in a Caucasian Australian population. *Pathology* 2005; 37 (2): 160 - 163.

Flett P, An update on the SA cerebral palsy register. *Action Packed*. Spring 2003.

Scher AI, Petterson B, Blair E, Ellenberg JH, Grether JK, Haan EA, Reddihough D, Yeargin-Allsopp M and Nelson KB. The risk of mortality or cerebral palsy in twins: A collaborative population-based study. *Pediatr Res* 2002;52: 671-81.

Gilbert M. What is cerebral palsy? *Nexus*, November, 2000.

Gilbert M. What is cerebral palsy and why have a register? *Action Packed*. Spring 1999.

Gilbert M. The South Australian cerebral palsy register. *Newsletter of the Australian Early Intervention Association (South Australian Chapter) Inc.* Winter, 1999.

Annual Reports

2003 Annual Report of the South Australian Cerebral Palsy Register.

2002 Annual Report of the South Australian Cerebral Palsy Register.

2001 Annual Report of the South Australian Cerebral Palsy Register.

2000 Annual Report of the South Australian Cerebral Palsy Register.

1999 Annual Report of the South Australian Cerebral Palsy Register.

Publications pertaining to cerebral palsy by members of the steering committee

Cerebral Palsy educational film for new families of children with cerebral palsy (work in progress and due to be completed August 2005).

Flett P, Review article, "Rehabilitation of spasticity and related problems in childhood cerebral palsy". *J Paed Child Health* 2003;39:6-14

Flett P. Book review: "Congenital hemiplegia", Neville B and Goodman R; eds., 2000. *J Paed Child Health* 2002;38:324.

Houltram J, Noble I, Boyd RN, Corry I, Flett P, Graham HK. Botulinum toxin type A in the management of equinus in children with cerebral palsy: an evidence-based economic evaluation. *Eur J Neurol* 2001; 8:S194-202.

Beal SM, Zeitz H, Connell T and Zschorn M. Athetoid quadriplegia and literacy. *J Paed Child Health* 2000; 36:389-391.

Flett P, Ferris R, McDonnell K, Raymond J, Walters G. Food for thought for people with disabilities and their carers. Ed. Sheedy C.T. Major revised edition. Project funded by Department of Human Services, Health Promotion SA, 1999.

Flett PJ, Stern LM, Waddy H, Connell TM, Seeger JD, Gibson SK. Botulinum toxin A versus fixed cast stretching for dynamic calf tightness in cerebral palsy. *J Paed Child Health* 1999; 35:71-77.

Appendix 1

GROSS MOTOR FUNCTION CLASSIFICATION SYSTEM IN CEREBRAL PALSY³

LEVEL I

Walks without restrictions: limitations in more advanced gross motor skills.

Before 2nd birthday: Infants move in and out of sitting and floor sit with both hands free to manipulate objects. Infants crawl on hands and knees, pull to stand and take steps holding onto furniture. Infants walk between 18 months and 2 years of age without the need for any assistive mobility device.

From age 2 to 4th birthday: Children floor sit with both hands free to manipulate objects. Movements in and out of floor sitting and standing are performed with adult assistance. Children walk as the preferred method of mobility without the need for any assistive mobility device.

From age 4 to 6th birthday: Children get into and out of and sit in a chair without the need for hand support. Children move from the floor and from chair sitting to standing without the need for objects for support. Children walk indoors and outdoors, and climb stairs. Emerging ability to run and jump.

From age 6 to 12: Children walk indoors and outdoors and climb stairs without limitations. Children perform gross motor skills including running and jumping but speed; balance and coordination are reduced.

LEVEL II

Walks without assistive devices: limitations walking outdoors and in the community.

Before 2nd birthday: Infants maintain floor sitting but may need to use their hands for support to maintain balance. Infants creep on their stomach or crawl on hands and knees. Infants may pull to stand and take steps holding onto furniture.

From age 2 to 4th birthday: Children floor sit but may have difficulty with balance when both hands are free to manipulate objects. Movements in and out of sitting are performed without adult assistance. Children pull to stand on a stable surface. Children crawl on hands and knees with a reciprocal pattern, cruise holding onto furniture and walk using an assistive mobility device as preferred methods of mobility.

From 4 to 6th birthday: Children sit in a chair with both hands free to manipulate objects. Children move from the floor to standing and from chair sitting to standing but often require a stable surface to push or pull up on with their arms. Children walk without the need for any assistive mobility device indoors and for short distances on level surfaces outdoors. Children climb stairs holding onto a railing but are unable to run or jump.

From age 6 to 12: Children walk indoors and outdoors and climb stairs holding onto a railing but experience limitations walking on uneven surfaces and inclines, and walking in crowds or confined spaces. Children have at best only minimal ability to perform gross motor skills such as running and jumping.

Distinction between Levels I and II

Compared with children in Level I, children in Level II have limitations in the ease of performing movement transitions: walking outdoors and in the community: the need for assistive mobility devices when beginning to walk: quality of movement: and the ability to perform gross motor skills such as running and jumping.

LEVEL III

Walks with assistive mobility devices: limitations walking outdoors and in the community.

Before 2nd birthday: Infants maintain floor sitting when the low back is supported. Infants roll and creep forward on their stomachs.

From age 2 to 4th birthday: Children maintain floor sitting often by 'W-sitting' (sitting between flexed and internally rotated hips and knees) and may require adult assistance to assume sitting. Children creep on their stomach or crawl on hands and knees (often without reciprocal leg movements) as their primary methods of self-mobility. Children may pull to stand on a stable surface and cruise short distances. Children may walk short distances indoors using an assistive mobility device and adult assistance for steering and turning.

From age 4 to 6th birthday: Children sit on a regular chair but may require pelvic or trunk support to maximise hand function. Children move in and out of chair sitting using a stable surface to push on or pull up with their arms. Children walk with an assistive mobility device on level surfaces and climb stairs with

Appendix 1 (continued)

assistance from an adult. Children frequently are transported when travelling for long distances or outdoors on uneven terrain.

From age 6 to 12: Children walk indoors or outdoors on a level surface with an assistive mobility device. Children may climb stairs holding onto a railing. Depending on upper limb function, children propel a wheelchair manually or are transported when travelling for long distances or outdoors on uneven terrain.

Distinction between Levels II and III

Differences are seen in the degree of achievement of functional mobility. Children in Level III need assistive mobility devices and frequently orthoses to walk, while children in Level II do not require assistive mobility devices after age 4.

LEVEL IV

Self-mobility with limitations: children are transported or use power mobility outdoors and in the community.

Before the 2nd birthday: Infants have head control but trunk support is required for floor sitting. Infants can roll to supine and may roll to prone.

From age 2 to 4th birthday: Children floor sit when placed, but are unable to maintain alignment and balance without use of their hands for support. Children frequently require adaptive equipment for sitting and standing. Self-mobility for short distances (within a room) is achieved through rolling, creeping on stomach, or crawling on hands and knees without reciprocal leg movement.

From age 4 to 6th birthday: Children sit on a chair but need adaptive seating for trunk control and to maximise hand function. Children move in and out of chair sitting with assistance from an adult or a stable surface to push or pull up on with their arms. Children may at best walk short distances with a walker and adult supervision but have difficulty turning and maintaining balance on uneven surfaces. Children are transported in the community. Children may achieve self-mobility using a power wheelchair.

From age 6 to 12: Children may maintain levels of function achieved before age 6 or rely more on wheeled mobility at home, school, and in the community. Children may achieve self-mobility using a power wheelchair.

Distinction between Levels III and IV

Differences in sitting ability and mobility exist, even allowing for extensive use of assistive technology. Children in Level III sit independently, have independent floor mobility and walk with assistive mobility devices. Children in Level IV function in sitting (usually supported) but independent mobility is very limited. Children in Level IV are more likely to be transported or use power mobility.

LEVEL V

Self-mobility is severely limited even with the use of assistive technology.

Before 2nd birthday: Physical impairments limit voluntary control of movement. Infants are unable to maintain antigravity head and trunk postures in prone and sitting. Infants require adult assistance to roll.

From age 2 to 12: Physical impairments restrict voluntary control of movement and the ability to maintain antigravity head and trunk postures. All areas of motor function are limited. Functional limitations in sitting and standing are not fully compensated for through the use of adaptive equipment and assistive technology. At Level V, children have no means of independent mobility and are transported. Some children achieve self-mobility using a power wheelchair with extensive adaptations.

Distinctions between Levels IV and V

Children in Level V lack independence even in basic antigravity postural control. Self-mobility is achieved only if the child can learn how to operate an electrically powered wheelchair. (Palisano et al, 1997)³.

Appendix 2

South Australian Cerebral Palsy Register Family Information Sheet

About the South Australian Cerebral Palsy Register

Approximately one in every 500 children born in South Australia has cerebral palsy. It is now recognised that most cerebral palsy is due to factors present before labour begins, and not as a result of events occurring during labour and delivery. However, what actually causes cerebral palsy is not clear. In order to determine these factors and to improve services, it is important to collect information about children with cerebral palsy.

For several years, doctors caring for children with cerebral palsy in South Australia discussed the possibility and value of establishing a cerebral palsy register. In 1991, a steering committee was formed which included doctors from Spastic Centres of SA, Novita Children's Services, Flinders Medical Centre, and the Women's and Children's Hospital. The South Australian Cerebral Palsy Register was established in 1997 with support from Novita Children's Services, Community Accommodation and Respite Agency (CARA), Rotary, and the Women's and Children's Hospital. The Register is located at, and receives its funding from the Women's and Children's Hospital.

What is the purpose of the Register?

The purpose of the Cerebral Palsy Register is to collect information about South Australian children with cerebral palsy. This information will enable us to:

- Find out how many children in South Australia have cerebral palsy;
- Detect changes in the number of children with cerebral palsy;
- Carry out research into the causes of cerebral palsy;
- Identify the full range of disabilities experienced by children with cerebral palsy;
- Help in the planning of services for children with cerebral palsy and;
- Increase knowledge in the community about cerebral palsy.

What benefits are there for my child?

At five years of age your child will receive a free, comprehensive clinical assessment by a specialist paediatrician. This assessment will take approximately 45 minutes. Information gained by the paediatrician may be used to make recommendations to your child's doctor about services or treatment. You will receive information of interest to you, including information about current research and the option to receive a copy of the Register's Annual Report.

What other benefits may come from participating?

By participating you will also benefit other children with cerebral palsy by contributing to research. Such research may:

- Find ways of preventing cerebral palsy;
- Find possible causes;
- Lead to new treatments or management strategies;
- Increase community awareness and recognition and;
- Lead to the development of new and/or improved services.

Who are the Register staff we will have contact with?

The person with whom you will have the most contact will be Kylie Tungaraza, the Coordinator of the South Australian Cerebral Palsy Register. Dr Peter Flett, Dr Ray Russo, Dr Andrew Tidemann, and Dr Deirdre White, are the specialist rehabilitation paediatricians who work with the Register. One of whom will conduct your child's clinical assessment.

Appendix 2 (continued)

How will information about my child be kept confidential?

All information contained on the South Australian Cerebral Palsy Register is strictly confidential. Only Register staff and those carrying out research in collaboration with them will use it. This means that no information, which identifies your child, will be released to other people unless you give written permission.

Are there any risks for my child?

There are no risks for your child associated with the medical assessment at five years of age. It will **not** involve the taking of blood samples or any other invasive tests.

Has the Register received approval from an Ethics Committee?

The South Australian Cerebral Palsy Register has received approval from the Research Ethics Committees of the Women's and Children's Hospital, Flinders Medical Centre, and the North Western Adelaide Health Service.

How do I give my consent?

You will be asked to give written consent to include your child on the Register. If, in the future, you change your mind, you can withdraw your consent. Refusing to include your child will not disadvantage your child in any way or change your relationship with health professionals or hospitals.

What happens now?

By now you will have received some information from your child's doctor about the South Australian Cerebral Palsy Register, and the Coordinator will have contacted you either by phone or letter.

The next step is to decide whether you would like your child to be included on the Register. If you decide that you would like your child to be included then you will be asked to sign a Consent Form and return it to the Coordinator of the Register. When your child is five years old, a paediatric rehabilitation specialist will conduct a medical assessment. This will usually be conducted at the Women's and Children's Hospital and will take approximately 45 minutes. It may be possible to make other arrangements for country families and those who would find it difficult to get to the Women's and Children's Hospital. The medical assessment will be conducted free of charge. You will receive no payment for your participation.

Although doctors working with the Cerebral Palsy Register will conduct the medical assessment, your child will remain in the care of their usual doctor. If, at any time, you have any questions about your child's treatment or development you should ask your child's usual doctor. Between your child's inclusion on the Register and their five year assessment, the Coordinator will maintain contact with you.

To collect complete information about your child's health it may be necessary to consult hospital records of your child's birth and current health (all of these options are explained on the Consent Form).

If you have any further questions please contact The Coordinator, South Australian Cerebral Palsy Register, on (08) 8161 7242 during office hours.

References

1. Badawi N, Watson L, Petterson B, Blair E, Slee J, Haan E, Stanley F. What constitutes cerebral palsy? *Developmental Medicine & Child Neurology* 1998; 40: 520-527.
2. Nelson KB, Dambrosia JM, Grether JK, Phillips TM. Neonatal cytokines and coagulation factors in children with cerebral palsy. *Annals of Neurology* 1998; 44: 665-675.
3. Palisano R, Rosenbaum P, Walter S, Russell D, Wood E, Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Developmental Medicine & Child Neurology* 1997; 39: 214-223.